



OPEN ACCESS

Estimating the information gap between emergency department records of community medication compared to on-line access to the community-based pharmacy records

Robyn Tamblyn,^{1,3} Lise Poissant,² Allen Huang,³ Nancy Winslade,³ Christian M Rochefort,^{3,4} Teresa Moraga,³ Pamela Doran³

¹Department of Epidemiology and Biostatistics, McGill University, Montreal, Canada

²Department of Rehabilitative Science, University of Montreal, Montreal, Canada

³Clinical and Health Informatics Research Group, Department of Medicine, McGill University, Montreal, Canada

⁴School of Nursing, McGill University, Montreal, Canada

Correspondence to

Dr Robyn Tamblyn, McGill University, Morrice House, 1140 Pine Ave W., Montreal, Quebec, Canada, H3A 1A3; robyn.tamblyn@mcgill.ca

Received 8 February 2013
Revised 28 May 2013
Accepted 24 June 2013
Published Online First
16 August 2013

ABSTRACT

Objective Errors in community medication histories increase the risk of adverse events. The objectives of this study were to estimate the extent to which access to community-based pharmacy records provided more information about prescription drug use than conventional medication histories.

Materials and methods A prospective cohort of patients with public drug insurance who visited the emergency departments (ED) in two teaching hospitals in Montreal, Quebec was recruited. Drug lists recorded in the patients' ED charts were compared with pharmacy records of dispensed medications retrieved from the public drug insurer. Patient and drug-related predictors of discrepancies were estimated using general estimating equation multivariate logistic regression.

Results 613 patients participated in the study (mean age 63.1 years, 59.2% women). Pharmacy records identified 41.5% more prescribed medications than were noted in the ED chart. Concordance was highest for anticoagulants, cardiovascular drugs and diuretics. Omissions in the ED chart were more common for drugs that may be taken episodically. Patients with more than 12 medications (OR 2.92, 95% CI 1.71 to 4.97) and more than one pharmacy (OR 3.85, 95% CI 1.80 to 6.59) were more likely to have omissions in the ED chart.

Discussion The development of health information exchanges could improve the efficiency and accuracy of information about community medication histories if they enable automated access to dispensed medication records from community pharmacies, particularly for the most vulnerable populations with multiple morbidities.

Conclusions Pharmacy records identified a substantial number of medications that were not in the ED chart. There is potential for greater safety and efficiency with automated access to pharmacy records.

BACKGROUND AND SIGNIFICANCE

Errors in the recording of prescription medication histories are distressingly common in the hospital setting, occurring in upwards of 50–70% of admitted patients.^{1–3} Over one quarter of these errors are attributable to incomplete medication histories obtained at the time of admission.⁴ In many cases, medication discrepancies lead to interrupted or inappropriate drug therapy during and after hospitalization,^{5–7} and are an important contributor to adverse drug events among hospitalized and recently discharged patients.^{3, 6} Following hospital

discharge, the perpetuation of these errors may result in unintended disruptions in essential treatment, drug interactions or therapeutic duplication—all of which have the potential to cause significant clinical harm and result in additional health-care costs.⁷ Indeed, a recent study suggests that there may be a 10% increase in adverse drug events among patients whose medication was unintentionally disrupted by a hospital stay.⁶

Although research in this area remains limited, some studies have shown significant discrepancies between medication lists generated during emergency department (ED) triage and those later verified by a pharmacist.^{1–3, 7, 8} The most common example is the omission of medications taken at home;^{2, 7} over 59% of medication histories recorded by physicians differ from those collected by pharmacy staff,² and although pharmacists are more effective in obtaining an accurate medication history, they are not available for all patient encounters.⁹ While computerized provider order entry systems are effective in minimizing the risk associated with prescribing errors, they cannot detect home medication use without linkage to a community pharmacy database.⁷ Better methods of obtaining the complete community drug list for each patient upon hospital admission would help to avoid preventable adverse drug events and increase the utility of community pharmacy systems in improving patient safety. The development of health information exchanges (HIE) in the USA and the pan-Canadian electronic health record in Canada are examples of technological infrastructures that could be used to provide access to community pharmacy records.^{10, 11} What remains unclear is whether access to community-based pharmacy records will provide substantially more information about prescription drug use than conventional medication histories.

Two recent studies, one conducted in Denmark and the other in the USA, set out to evaluate the magnitude of discrepancy between these two sources of information; however, their results were conflicting. In the Danish study, 27% of medications documented in electronic pharmacy records were not reported by patients on admission to an acute medical emergency ward, whereas only 6% of medications reported by patients were omitted from pharmacy records.¹² In contrast, in the American study, 30% of medications reported by patients were not included in an insurance claims



Open Access
Scan to access more
free content

To cite: Tamblyn R, Poissant L, Huang A, et al. *J Am Med Inform Assoc* 2014;**21**:391–398.

Research and applications

database, possibly because unlike pharmacy records, which record all prescriptions dispensed, insurance claims only include those drugs that are covered.¹³ Neither of these studies identified the characteristics of patients at risk of having discrepancies in their medication histories. These data are needed to evaluate the extent to which investment in and maintenance of HIE will produce clinical value for patients.

OBJECTIVE

The purpose of this study was to determine the extent to which community medication histories documented in the ED differed from records of dispensed medications provided by community pharmacies, and the patient and drug-related predictors of discrepancies that could be used to identify higher risk groups.

MATERIALS AND METHODS

Setting

This study took place at two teaching hospitals in Quebec, Canada, where all residents have provincial medical insurance, and approximately 50% of the population is covered by public drug insurance. Claims records for all dispensed medications from Quebec pharmacies and medical services from fee-for-service physicians are maintained at the Régie de l'Assurance Maladie du Québec (RAMQ) and can be linked using a unique lifetime identifier to create a longitudinal health record for each provincial resident.

Study design and participants

A prospective cohort study was conducted. Individuals who were seen in the ED at the Royal Victoria Hospital and the Montreal General Hospital between 21 May and 27 August 2008, who were 18 years of age or older, reported that they took at least one prescription medication and who had public drug insurance were asked to participate. Research assistants assessed eligibility and sought patient consent. The McGill University Health Centre's ethics committee, the RAMQ, and the provincial privacy commission approved the study.

Data collection and retrieval

For each consenting patient, two different community drug lists were retrieved: one that contained the drug list recorded by the triage nurse and ED staff in the patient's paper-based medical chart at the time of their ED visit, and another obtained from the community pharmacy records of dispensed medications recorded in the administrative databases of the RAMQ. For patients with multiple ED visits during the study period, only information pertaining to their first visit was collected from the ED chart and the community pharmacy records.

Chart retrieval

Trained research assistants reviewed all prescription drugs recorded in the ED section of the patient's hospital chart and entered the community drug list (drug name, strength, form) documented at the time of the visit into an electronic database. To minimize data entry errors, the database auto-generated a list of possible drug names from the first three typed letters; once the drug was selected, the strength and form was selected from a drop-down list, or entered in a free text field if the correct option was not available. If a drug name, strength, or form was missing or illegible, this information was also recorded in the database. All drugs, both prescription and over the counter, were coded by drug identification number, when possible, and/or by the generic drug code (ie, by drug ingredient regardless of manufacturer or format) using a proprietorial classification

system (Vigilance Santé).¹⁴ The number of illegible drugs was recorded for each patient, when applicable.

Provincial insurer (RAMQ) medication list retrieval from Quebec pharmacies

For each consenting patient, data on all prescription drugs dispensed from Quebec pharmacies in the year before the ED visit date were retrieved from the RAMQ. Data included medication drug identification number, strength, format, quantity dispensed, the duration of the prescription, and encrypted identifiers for each of the patients, prescribing physician, and dispensing pharmacy. Dispensed drugs were grouped by generic code and those with an active prescription within the 2 months before the ED visit date were classified as current community medications. Patients using medications for chronic conditions may miss 20% of doses (a mean compliance of 80%), and thus a window of 2 months back from the ED visit date was used to identify current medication.

Main outcome

The primary outcome was a discrepancy between community medications that were documented in the ED chart compared to dispensed medication records from the RAMQ databases. A discrepancy was defined as the absence of a drug with the same generic drug code in either the ED or RAMQ drug lists. All illegible drug entries in the ED chart were excluded.

Predictors of discrepancies

Potential patient and drug-related predictors of medication list discrepancies were measured with the aim of identifying subsets of patients who may be at greater risk of an incomplete community medication history. As many jurisdictions do not have access to records of dispensed medications, we assessed patient characteristics that were associated with a greater number of medications in pharmacy-recorded claims relative to the ED chart. Using a combination of medical chart and administrative data, we measured patient age and sex, health conditions that would create potential communication barriers,¹⁵ as well as the number and therapeutic classes of medications as these factors may influence patient recall and communication. Using administrative data we measured the frequency of ED visits, and hospitalization in the past year, as well as the number of pharmacies and physicians involved in prescribing and dispensing as these factors create challenges in the continuity of care. Published algorithms were employed to measure patient and healthcare utilization characteristics.^{16–19} Over-the-counter drugs were excluded, as they are not recorded in the pharmacy records.

Statistical analysis

Descriptive statistics were used to characterize the cohort, and to examine the frequency of discrepancies between the ED and Quebec pharmacy dispensing records for prescription drugs, by class and most frequent generic drugs. Multivariate logistic regression was used to estimate the association between patient demographics, health conditions, and healthcare utilization, as well as the likelihood of having more prescribed drugs listed in the pharmacy records than the ED chart. We estimated both the univariate and multivariate associations for each variable. To avoid multi-collinearity, we selected only one variable when multiple variables measuring the same concept were used (eg, number and types of drugs). SAS V.9.2 was used for all analyses.

RESULTS

Overall, 613 patients met the eligibility criteria and were enrolled in the study. The mean patient age was 63.1 years, 59.2% were women, and 32.6% had health conditions that may have created potential communication barriers (table 1). In the year before the ED visit, 45.7% of patients had four or more prescribing physicians, and 7.5% had four or more pharmacies.

Approximately one quarter (26.1%) of patients had more than 12 active prescription drugs documented at the ED visit, either in the hospital chart or through the pharmacy records; 57.3% used at least one over-the-counter medication, and 14.5% had at least one illegible drug in the ED chart (table 2).

The most prevalent over-the-counter medications were acetaminophen, acetylsalicylic acid, multivitamins, and calcium (table 3). For prescribed medications, 30.8% of patients had a single prescribing physician, and 72.4% had a single dispensing pharmacy. Cardiovascular drugs, central nervous system agents,

Table 1 Demographic and healthcare utilization characteristics of the 613 patients in the study population in the year before the ED visit

| Demographic characteristics | Mean | SD |
|--|------|------|
| Patient age (years) | 63.1 | 18.8 |
| | N | % |
| <50 years | 150 | 24.5 |
| 50–70 years | 197 | 32.1 |
| >70 years | 266 | 43.4 |
| Sex | | |
| Female | 363 | 59.2 |
| Male | 250 | 40.8 |
| Health conditions with communication barriers | | |
| Any condition | 200 | 32.6 |
| Psychiatric disorder* | 183 | 29.9 |
| Social distancing problem† | 32 | 5.2 |
| Physical communication problems‡ | 3 | 0.5 |
| Healthcare utilization in past year | | |
| No of ED visits in past year | | |
| No visits | 307 | 50.1 |
| 1 Visit | 141 | 23.0 |
| 2–3 Visits | 104 | 17.0 |
| ≥4 Visits | 61 | 10.0 |
| No of hospitalizations in past year | | |
| No hospitalizations | 412 | 67.2 |
| 1 Hospitalization | 116 | 18.9 |
| ≥2 Hospitalizations | 85 | 13.9 |
| Continuity of medication management in past year | | |
| No of prescribing physicians in past year | | |
| No information on prescribers | 35 | 5.7 |
| 1 Prescriber | 84 | 13.7 |
| 2–3 Prescribers | 214 | 34.9 |
| ≥4 Prescribers | 280 | 45.7 |
| No of pharmacies in past year | | |
| No information on pharmacies | 33 | 4.5 |
| 1 Pharmacy | 327 | 53.3 |
| 2–3 Pharmacies | 207 | 33.8 |
| ≥4 Pharmacies | 46 | 7.5 |

*Including diagnoses for depression, anxiety and adjustment disorders, schizophrenia and other psychotic disorders, mood disorder including bipolar and personality disorder.

†Includes alcoholism, drug abuse and homelessness.

‡Includes documented communication difficulties, blindness and deafness. ED, emergency department.

Table 2 Characteristics of prescribed and OTC medication use at ED visit and discrepancies between ED and pharmacy record sources at the time of the ED visit

| No of prescribed drugs (either ED or pharmacy source) | N | % |
|--|-----|------|
| 0 Drugs (only OTC) | 8 | 1.3 |
| 1 Drug | 36 | 5.9 |
| 2–6 Drugs | 220 | 35.9 |
| 7–12 Drugs | 189 | 30.8 |
| >12 Drugs | 160 | 26.1 |
| Patients with ≥1 OTC medication | 351 | 57.3 |
| Patients with ≥1 illegible drug in the ED chart | 89 | 14.5 |
| Drug use by therapy class | | |
| Central nervous system agents | 360 | 58.7 |
| Cardiovascular drugs | 363 | 59.2 |
| Hormones and synthetic substitutes | 282 | 46.0 |
| Blood formation, coagulation and thrombosis | 218 | 35.5 |
| Gastrointestinal drugs | 211 | 34.4 |
| Electrolytic, caloric and water balance | 237 | 38.9 |
| Anti-infective agents | 162 | 26.4 |
| Autonomic drugs | 161 | 26.3 |
| Eye, ear, nose and throat preparations | 93 | 15.7 |
| Other | 273 | 44.5 |
| No of prescribing physicians | | |
| No information on prescribers | 45 | 7.3 |
| 1 Prescriber | 189 | 30.8 |
| 2–3 Prescribers | 265 | 43.2 |
| ≥4 Prescribers | 114 | 18.6 |
| No of pharmacies | | |
| No information on pharmacies | 45 | 7.3 |
| 1 Pharmacy | 444 | 72.4 |
| 2–3 Pharmacies | 114 | 18.6 |
| ≥4 Pharmacies | 10 | 1.6 |
| Differences between pharmacy records and ED chart | | |
| Pharmacy records document more prescribed drugs than ED chart | 463 | 75.5 |
| 1 More drug | 122 | 19.9 |
| 2–4 More drugs | 219 | 35.7 |
| ≥5 Drugs | 122 | 19.9 |
| ED chart documents more prescribed drugs than pharmacy records | 73 | 11.9 |
| 1 More drug | 31 | 5.0 |
| 2–4 More drugs | 25 | 4.1 |
| ≥5 Drugs | 17 | 2.8 |

ED, emergency department; OTC, over-the-counter.

hormones and synthetic substitutes were the most prevalent therapies Pharmacy records documented more active drugs than the ED chart in 75.5% of patients, and in 11.9%, the ED chart listed more medications.

There was concordance between the ED chart and pharmacy records for 45.2% of medications, 41.5% were only identified in the pharmacy records, and 13.4% were only in the ED chart. Concordance was highest for anticoagulants (64.6%), cardiovascular drugs (61.4%), and diuretics (56.2%) (table 4).

Drugs that are taken episodically were more frequently recorded in pharmacy records only (table 4). For example, 80.9% of skin agents, 75.3% of anti-infective drugs, 68.6% of eye, ear nose and throat medications, 46.2% of central nervous system agents, 44.0% of gastrointestinal drugs, and 43.4% of autonomic drugs, such as bronchodilators, were only recorded in pharmacy records. In the case of anti-infective drugs, 91% of patients who used anti-infective agents had a prescription end-date that preceded the ED visit; however, 34% of patients

Research and applications

Table 3 Frequency distribution of OTC drugs documented in the ED chart by therapeutic class and most frequently used drugs

| Therapeutic class | N | % of all OTC drugs | % of patients using drug |
|---|-----|--------------------|--------------------------|
| Central nervous system agents | 175 | 24.8 | 28.5 |
| Acetaminophen | 85 | 12.1 | 13.9 |
| ASA | 80 | 11.3 | 13.1 |
| Ibuprofen | 9 | 1.3 | 1.5 |
| Electrolytic, caloric and water balance | 118 | 16.7 | 19.2 |
| Calcium carbonate+vitamin D | 35 | 5.0 | 5.7 |
| Calcium carbonate | 28 | 4.0 | 4.6 |
| Calcium citrate | 28 | 4.0 | 4.6 |
| Blood formation, coagulation and thrombosis | 111 | 15.7 | 18.1 |
| ASA antiplatelet | 85 | 12.1 | 13.9 |
| Ferrous sulfate | 12 | 1.7 | 2.0 |
| Iron-chelated | 7 | 1.0 | 1.1 |
| Vitamins | 95 | 13.5 | 15.5 |
| Multivitamins | 39 | 5.5 | 6.4 |
| Vitamin D3 | 18 | 2.6 | 2.9 |
| Vitamin D | 12 | 1.1 | 2.0 |
| Gastrointestinal drugs | 101 | 14.3 | 16.5 |
| Docusate sodium | 38 | 5.4 | 6.2 |
| Sennosides A–B | 21 | 3.0 | 3.4 |
| Dimenhydrinate | 9 | 1.3 | 1.5 |
| Skin and mucous membrane agents | 27 | 3.8 | 4.4 |
| Petrolatum-white | 8 | 1.1 | 1.3 |
| Gramicidin+PO Lymyxin B | 3 | 0.4 | 0.5 |
| Clotrimazole | 3 | 0.4 | 0.5 |
| Other* | 87 | 11.4 | 14.1 |
| Quinine sulfate | 9 | 1.3 | 1.5 |
| Potassium iodide+ammonium chloride | 7 | 1.0 | 1.1 |
| Products natural miscellaneous | 6 | 0.9 | 1.0 |

*Other: Miscellaneous therapeutic agents, antihistamine drugs, respiratory tract agents, diagnostic agents, devices, anti-infective agents, autonomic drugs, eye, ear, nose and throat preparations.
ASA, acetylsalicylic acid; ED, emergency department; OTC, over-the-counter.

were prescribed multiple different anti-infective agents in the past 2 months. Of interest, 9.9% (gastrointestinal drugs) to 18.7% (autonomic drugs) of drugs were listed in the ED chart but were not found in pharmacy records of dispensed medications, at least in the 2 months before the ED visit.

The likelihood of having more drugs documented in pharmacy records than the ED chart was associated with the number of prescribed drugs, and the number of pharmacies, but not age, sex, hospitalizations or communication problems. (table 5). Patients with more frequent ED visits were more likely to have discrepancies in bivariate analysis, but these trends were largely explained by the number of medications and healthcare use and were not significant in multivariate analysis.

Patients using more than 12 drugs were almost three times more likely (OR 2.92, 95% CI 1.71 to 4.97) to have drugs documented in pharmacy records that were not noted in the ED chart. Compared to cardiovascular drugs, anti-infective drugs, blood formation, coagulation and thrombosis drugs, gastrointestinal drugs, and central nervous system drugs were more likely to be documented only in pharmacy records and not the ED chart. The number of prescribing physicians, both in the past year and for current medications, was associated with the

likelihood of having more drugs documented in pharmacy records. However, the number of prescribing physicians was also strongly correlated with the number of drugs ($r=0.54$, $p<.0001$), so only the number of drugs was included in the multivariate models. Using more than one pharmacy, both in the past year and for current medication, was associated with more than a threefold increase (OR 3.45, 95% CI 1.80 to 6.59) in the risk of having more medications listed in pharmacy records, even after adjusting for the number of current medications.

DISCUSSION

This study provides insight into the potential value of accessing information electronically from community pharmacy medication records at the time of an ED visit. Pharmacy records identified 41.5% more prescribed medications that were not noted in the ED chart. More vulnerable patients, that is, those with many medications, many prescribing physicians, and multiple ED visits were more likely to have omissions in their ED record of current medications. Although having potential communication problems did not turn out to be a significant predictor of medication discrepancies, patients who were unconscious or agitated would have been less likely to be enrolled in this study because of known consent biases,²⁰ and thus we probably underestimated these effects. We found that omissions were more common for some groups of medications than others, particularly for therapeutic classes of drugs that may be taken episodically to control symptoms: anxiolytics, analgesics, gastrointestinal drugs, and asthma medications. The clinical significance of discontinuing these medications, even when intended, is substantial as discontinuation results in an adverse event rate of 14.3% (gastrointestinal drugs) to 29.7% (anxiolytics) in older people.²¹ These groups of medications are more likely to be 'forgotten' by patients in self-reported surveys of medication use^{7 22–24} and ED medication histories² when compared to home-based assessment of medicine cabinets,^{25 26} and pharmacist review.² Although most cardiovascular and anticoagulant medications were documented in both the ED chart and pharmacy records, 29.2% of patients prescribed clopidogrel, and 8.3% of those prescribed warfarin had these medications only documented in pharmacy records. Although ED staff may selectively solicit information about these medications (particularly anticoagulants) as these drugs are frequently involved in adverse drug events,^{27–30} there is clearly benefit in obtaining pharmacy records for patients on these medications.

Although pharmacy records of dispensed medications do not reflect actual usage, they can provide a clinically useful proxy. Several validation studies have shown that medication possession ratios calculated from pharmacy records are predictive of various measures of disease control such as blood pressure, glycosylated hemoglobin levels, and asthma exacerbation.^{31–35} There is also excellent agreement between dispensing records and the use of antihypertensive drugs.^{31 32 36 37} Moreover, if access to dispensed medications records from community pharmacies is automated, there is considerable potential to improve efficiencies for hospital-based pharmacists in documenting the best medication history,^{38 39} as the average time spent per patient for this task can range from 13 to 90 min.^{7 40 41}

To optimize the utility of access to computerized community-based pharmacy records, a few issues need to be addressed. First, a decision needs to be made about the time window that will be used to retrieve data. In this study, we included medications in the 2 months before the ED visit. For patients with poor compliance, we found that we could only pick up

Table 4 Discrepancies in prescribed drugs between pharmacy records* and the ED chart by therapeutic class

| Therapeutic class† | Total | ED only (n, %) | Pharmacy only (n, %) | Documented in both (n, %) |
|---|-------|----------------|----------------------|---------------------------|
| Cardiovascular drugs | 903 | 113 (12.5) | 236 (26.1) | 554 (61.4) |
| Atorvastatin | 120 | 13 (10.8) | 27 (22.5) | 80 (66.7) |
| Amlodipine | 80 | 3 (3.8) | 15 (18.8) | 62 (75.5) |
| Metoprolol | 69 | 8 (11.6) | 12 (17.4) | 49 (71.0) |
| Central nervous system agents | 716 | 121 (16.9) | 331 (46.2) | 264 (36.9) |
| Lorazepam | 64 | 8 (12.5) | 27 (42.2) | 29 (45.3) |
| Naproxen | 42 | 6 (14.3) | 28 (66.7) | 8 (19.0) |
| Codeine+acetaminophen | 40 | 6 (15.0) | 30 (75.0) | 4 (10.0) |
| Hormones and synthetic substitutes | 408 | 47 (11.5) | 134 (32.8) | 227 (55.6) |
| Levothyroxine | 97 | 5 (5.2) | 18 (18.6) | 74 (76.3) |
| Metformin | 87 | 7 (8.0) | 9 (10.3) | 71 (81.6) |
| Prednisone | 38 | 2 (5.3) | 20 (52.6) | 16 (42.1) |
| Blood formation, coagulation and thrombosis | 96 | 15 (15.6) | 19 (19.8) | 62 (64.6) |
| Warfarin | 48 | 9 (18.8) | 4 (8.3) | 35 (72.9) |
| Clopidogrel | 24 | 1 (4.2) | 7 (29.2) | 16 (66.7) |
| Dalteparin | 4 | 1 (25.0) | 2 (50.0) | 1 (25.0) |
| Gastrointestinal drugs | 232 | 23 (9.9) | 102 (44.0) | 107 (46.1) |
| Pantoprazole | 108 | 13 (12.0) | 43 (39.8) | 52 (48.1) |
| Esomeprazole | 33 | 5 (15.2) | 14 (42.4) | 14 (42.4) |
| Omeprazole | 29 | 0 | 13 (44.8) | 16 (55.2) |
| Electrolytic, caloric and water balance | 169 | 17 (10.1) | 57 (33.7) | 95 (56.2) |
| Furosemide | 65 | 8 (12.3) | 17 (26.6) | 40 (61.5) |
| Hydrochlorothiazide | 65 | 8 (12.3) | 20 (30.8) | 37 (56.9) |
| Indapamide | 16 | 1 (6.3) | 7 (43.8) | 8 (50.0) |
| Anti-infective agents | 239 | 24 (10.0) | 180 (75.3) | 35 (14.6) |
| Ciprofloxacin | 36 | 3 (8.3) | 28 (77.8) | 5 (13.9) |
| Amoxicillin | 25 | 2 (8.0) | 20 (80.0) | 3 (12.0) |
| Azithromycin | 18 | 2 (11.1) | 16 (88.9) | 0 |
| Autonomic drugs | 249 | 49 (18.7) | 108 (43.4) | 92 (36.9) |
| Salbutamol | 74 | 19 (25.7) | 37 (50.0) | 18 (24.3) |
| Tamsulosin | 31 | 5 (16.1) | 10 (32.3) | 16 (51.6) |
| Tiotropium | 26 | 4 (15.4) | 11 (42.3) | 11 (42.3) |
| Eye, ear, nose and throat preparations | 172 | 25 (14.5) | 118 (68.6) | 29 (16.9) |
| Fluticasone | 53 | 8 (15.1) | 36 (67.9) | 9 (17.0) |
| Latanoprost | 16 | 2 (12.5) | 10 (62.5) | 4 (25.0) |
| Mometasone | 16 | 3 (18.8) | 12 (75.0) | 1 (6.3) |
| Skin and mucous membrane agents | 89 | 10 (11.2) | 72 (80.9) | 7 (7.9) |
| Betamethasone valerate | 8 | 1 (12.5) | 7 (87.5) | 0 |
| Mometasone furoate | 8 | 1 (12.5) | 7 (87.5) | 0 |
| Fluocinonide | 6 | 1 (16.7) | 5 (83.3) | 0 |
| Other‡ | 208 | 22 (10.6) | 86 (41.3) | 100 (48.1) |
| Alendronate | 31 | 2 (6.5) | 12 (38.7) | 17 (54.8) |
| Risedronate | 30 | 3 (10.0) | 11 (36.7) | 16 (53.3) |
| Allopurinol | 18 | 1 (5.6) | 5 (27.8) | 12 (66.7) |
| Overall | 3481 | 466 (13.4) | 1443 (41.5) | 1572 (45.2) |

*Active prescribed drugs in the 2 months before the ED visit.

†Includes the three most common drugs in each therapeutic class.

‡Including respiratory tract agents, devices, antineoplastic agents, miscellaneous therapeutic agents, diagnostic agents, vitamins, smooth and muscle relaxants, heavy metal antagonists. ED, emergency department.

medications that were only documented in the ED chart if we looked back 12 months before the ED visit. In contrast, a shorter time window may be more appropriate to exclude short-term treatments such as antibiotics, which were likely to be 'over-reported' as current medications, because 91% had an end-date that preceded the ED visit. However, of interest was the fact that we found that 34% of patients were taking more than one anti-infective drug in the past 2 months. This information could be relevant to ED staff in detecting adverse anti-

infective drug events or in avoiding previous failed treatment choices. To accommodate these issues, the window for retrieving medications for clinical use needs to be adjustable, allowing the user to conduct reviews by time period, therapy class, population, and insurance cost-sharing policies that affect compliance. Mean prescription duration is also an important factor to consider, as in some jurisdictions 3-month durations (versus 1 month in Quebec) are the norm, and thus a longer retrieval window will be needed.

Research and applications

Table 5 Factors associated with discrepancies: more drugs in pharmacy records than documented in the ED chart

| Factors | More drugs pharmacy records N (%) | Bivariate analysis | | Multivariate analysis | |
|---------------------------------|--------------------------------------|-----------------------|---------|------------------------|---------|
| | | OR (95% CI) | p Value | OR (95% CI) | p Value |
| Patient age (years) | | | | | |
| <50 | 111 (23.9) | Ref. | | Ref. | |
| 50–70 | 149 (32.0) | 1.09 (0.67 to 1.78) | 0.72 | 0.875 (0.504 to 1.518) | 0.64 |
| >70 | 205 (44.1) | 1.18 (0.75 to 1.88) | 0.47 | 0.814 (0.470 to 1.412) | 0.46 |
| Gender | | | | | |
| Female | 273 (58.7) | Ref. | | | |
| Male | 192 (41.3) | 1.09 (0.75 to 1.59) | 0.66 | 1.00 (0.67 to 1.49) | 0.99 |
| Communication barrier | 158 (34.0) | 1.29 (0.86 to 1.94) | 0.22 | 1.08 (0.69 to 1.67) | 0.75 |
| No of prescribed drugs | | | | | |
| 1 Drug | 18 (3.9) | 0.66 (0.31 to 1.42) | 0.29 | 0.74 (0.33 to 1.68) | 0.47 |
| 2–6 Drugs | 137 (29.5) | Ref. | | Ref. | |
| 7–12 Drugs | 121 (26.0) | 1.86 (1.16 to 3.00) | 0.01 | 1.99 (1.21 to 3.29) | 0.007 |
| ≥12 Drugs | 166 (33.3) | 2.92 (1.80 to 4.76) | <.0001 | 2.92 (1.71 to 4.97) | <.0001 |
| Therapeutic class† | | | | | |
| Cardiovascular | 286 (61.5) | Ref. | | | |
| Central nervous system | 289 (62.2) | 1.64 (1.09 to 2.46) | 0.01 | | |
| Hormones/synthetic subs. | 222 (47.7) | 1.11 (0.73 to 1.67) | 0.63 | | |
| Blood, coag/thrombosis | 192 (41.3) | 2.96 (1.79 to 4.89) | <.0001 | | |
| Gastrointestinal | 181 (38.9) | 1.69 (1.05 to 2.71) | <.0001 | | |
| Electrolytic, caloric and water | 196 (42.2) | 1.12 (0.71 to 1.76) | 0.64 | | |
| Anti-infectives | 145 (31.2) | 3.49 (1.99 to 6.12) | <0.001 | | |
| Autonomic | 128 (27.5) | 0.72 (0.44 to 1.19) | 0.20 | | |
| Eye, ear, nose and throat | 77 (16.6) | 0.96 (0.51 to 1.79) | 0.89 | | |
| Other | 236 (50.8) | 2.46 (1.57 to 3.86) | <.0001 | | |
| ED visits in past year | | | | | |
| No visits | 218 (46.9) | Ref. | | Ref. | |
| 1 Visit | 107 (23.0) | 1.28 (0.81 to 2.02) | 0.30 | 1.14 (0.70 to 1.84) | 0.61 |
| 2–3 Visits | 85 (18.3) | 1.80 (1.03 to 3.12) | 0.04 | 1.58 (0.86 to 2.89) | 0.14 |
| ≥4 Visits | 55 (11.8) | 3.50 (1.49 to 8.23) | 0.004 | 2.33 (0.91 to 5.96) | 0.08 |
| Hospitalizations in past year | | | | | |
| No hospitalization | 304 (65.4) | Ref. | | Ref. | |
| ≥1 Hospitalization | 161 (34.6) | 1.42 (0.95 to 2.14) | 0.09 | 0.93 (0.58 to 1.50) | 0.76 |
| Current pharmacies | | | | | |
| 1 Pharmacy | 352 (75.7) | Ref. | | Ref. | |
| ≥2 Pharmacies | 113 (24.3) | 3.85 (2.03 to 7.30) | <.0001 | 3.45 (1.80 to 6.59) | 0.0002 |
| Current prescribing MD* | | | | | |
| 1 Prescriber | 128 (27.5) | Ref. | | | |
| 2–3 Prescribers | 228 (49.0) | 5.05 (3.28 to 7.77) | <.0001 | | |
| ≥4 Prescribers | 109 (23.4) | 16.50 (6.72 to 40.50) | <.0001 | | |
| Pharmacies past year† | | | | | |
| 1 Pharmacy | 253 (54.4) | Ref. | | | |
| 2–3 Pharmacies | 172 (37.0) | 2.06 (1.35 to 3.16) | <.0001 | | |
| ≥4 Pharmacies | 40 (8.6) | 2.64 (1.11 to 6.23) | 0.03 | | |
| Prescribing MD past year* | | | | | |
| 1 Prescriber | 55 (11.8) | Ref. | | | |
| 2–3 Prescribers | 161 (34.6) | 3.50 (2.18 to 5.64) | <.0001 | | |
| ≥4 Prescribers | 249 (53.5) | 9.21 (5.45 to 15.44) | <.0001 | | |

*Number of prescribing physicians and number of drugs was strongly correlated ($r=0.54$) so only number of drugs was included in the model.

†Therapeutic drug classes and number of drugs was correlated so only number of drugs was included in the multivariate model. We selected the utilization variable for number of pharmacies that was most strongly associated with the outcome.
ED, emergency department; MD, physician.

Our results demonstrate the potential value in having access to community-based pharmacy records through infrastructures such as HIE, because they provide more information about prescription drug use compared to conventional medication histories. However, the cost of integrating and sustaining electronic access to these records would need to be justified by either a

reduction in professional time for medication history-taking (which is a substantial barrier to the medication reconciliation process), and/or in preventable adverse drug events. To date, cost–benefit models of HIE have been largely theoretical, based on economic modeling of assumed benefits and cost-savings.^{42 43} The lack of empirical evidence of the sources and types of cost

saving may be one reason why most regional HIE in the USA have not established viable business models for financial sustainability.⁴⁴ A recent evaluation of HIE in ED care in Tennessee is among one of the first studies to provide evidence that HIE implementation can reduce hospitalizations, and in some cases, can reduce diagnostic and laboratory test ordering, producing an annual cost savings of US\$1.07 million.⁴⁵

Greater efficiencies and cost savings in professional time could be gained with automated access to pharmacy records if patients were provided with an opportunity to verify the medications they were taking. An interesting approach was evaluated in the Veteran's hospital system. Kiosks were available in the ED for patients to retrieve and verify medications they were actually taking based on records of dispensed prescriptions.⁴⁶ A majority of the ED staff agreed that the tool was effective; however, the overwhelming volume of information made identifying discrepancies in the medication lists difficult, and some providers felt that it increased their workload. Despite these shortcomings, the kiosk was successfully integrated in the ED workflow and even patients with poor computer skills were able to use it. Similar features could be included in personal health records and patient portals that are linked to community pharmacy information systems, and this may be a fruitful area for future research.

Limitations that need to be considered are that the study was based in two teaching hospitals where electronic health records were not available. The benefits of automating access to community pharmacy records may be greater for hospitals that are still using paper-based processes as we found that 14.5% of patients had at least one illegible drug in their ED chart, highlighting one of the many benefits of electronic documentation of medication histories. Only publicly insured patients were included, and this subpopulation may be more vulnerable as it includes both older patients and those receiving social assistance. We did not have a definitive 'gold standard' of the current medication list by which the accuracy of the ED list, or the additional drugs identified from community pharmacies could be judged. Moreover, as drug doses and routes were rarely recorded in the chart, we probably underestimated discrepancies related to dose and route.

CONCLUSION

In summary, we found community pharmacy dispensed medication records identified a substantial number of additional medications that were not noted in the ED chart, particularly for the most vulnerable patients. There is potential to gain greater safety and efficiency within hospital ED with automated access to these pharmacy records.

Contributors RT, LP, AH, NW and CMR contributed to the conception and design, acquisition of data, or analysis and interpretation of data, and drafted the article or critically revised it for important intellectual content. TM completed the data analysis and PD contributed to drafting and revising the article. All authors approved the final version to be published.

Funding Funding was provided by the: Canadian Institutes of Health Research.

Competing interests None.

Ethics approval The McGill University Health Centre's ethics committee, the RAMQ, and the provincial privacy commission approved the study.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

REFERENCES

- Andersen SE, Pedersen AB, Bach KF. Medication history on internal medicine wards: assessment of extra information collected from second drug interviews and GP lists. *Pharmacoepidemiol Drug Saf* 2003;12:491–8.
- de Winter S, Spriet I, Indevuyst C, et al. Pharmacist- versus physician-acquired medication history: a prospective study at the emergency department. *Qual Saf Health Care* 2010;19:371–5.
- Slain D, Kincaid SE, Dunswoth TS. Discrepancies between home medications listed at hospital admission and reported medical conditions. *Am J Geriatr Pharmacother* 2008;6:161–6.
- Tam VC, Knowles SR, Cornish PL, et al. Frequency, type and clinical importance of medication history errors at admission to hospital: a systematic review. *CMAJ* 2005;173:510–15.
- Bell CM, Bajcar J, Bierman AS, et al. Potentially unintended discontinuation of long-term medication use after elective surgical procedures. *Arch Intern Med* 2006;166:2525–31.
- Bell CM, Brener SS, Gunraj N, et al. Association of ICU or hospital admission with unintentional discontinuation of medications for chronic diseases. *JAMA* 2011;306:840–7.
- Cornish PL, Knowles SR, Marchesano R, et al. Unintended medication discrepancies at the time of hospital admission. *Arch Intern Med* 2005;165:424–9.
- Mazer P, Gischler SJ, Koot HM, et al. Impact of a child with congenital anomalies on parents (ICCAP) questionnaire; a psychometric analysis. *Health Qual Life Outcomes* 2008;6:102;2.
- Mersfelder TL, Bickel RJ. Inpatient medication history verification by pharmacy students. *Am J Health Syst Pharm* 2008;65:2273–5.
- Webster PC. Centralized, nationwide electronic health records schemes under assault. *CMAJ* 2011;183:E1105–6.
- Kuperman GJ. Health-information exchange: why are we doing it, and what are we doing? *J Am Med Inform Assoc* 2011;18:678–82.
- Glintborg B, Poulsen HE, Dalhoff KP. The use of nationwide on-line prescription records improves the drug history in hospitalized patients. *Br J Clin Pharmacol* 2007;65:265–9.
- Warholak TL, McCulloch M, Baumgart A, et al. An exploratory comparison of medication lists at hospital admission with administrative database records. *JMCP* 2009;15:751.
- Vigilance Santé, Quebec, Canada. 2013. <http://www.vigilance.ca/>.
- Bartlett G, Blais R, Tamblin R, et al. Impact of patient communication problems on the risk of preventable adverse events in acute care settings. *CMAJ* 2008;178:1555–62.
- Tamblin R, Reid T, Mayo N, et al. Using medical services claims to assess injuries in the elderly: sensitivity of diagnostic and procedure codes for injury ascertainment. *J Clin Epidemiol* 2000;53:183–94.
- Tamblin R, Laprise R, Hanley JA, et al. Adverse events associated with prescription drug cost-sharing among poor and elderly persons. *JAMA* 2001;285:421–9.
- Tamblin R, Abrahamowicz M, Dauphinee WD, et al. Association between licensure examination scores and practice in primary care. *JAMA* 2002;288:3019–26.
- Wilchesky M, Tamblin RM, Huang A. Validation of diagnostic codes within medical services claims. *J Clin Epidemiol* 2004;57:131–41.
- Tu JV, Willison DJ, Silver FL, et al. Impracticability of informed consent in the Registry of the Canadian Stroke Network. *N Engl J Med* 2004;350:1414–21.
- Graves T, Hanlon JT, Schmader KE, et al. Adverse events after discontinuing medications in elderly outpatients. *Arch Intern Med* 1997;157:2205–10.
- Lau HS, Florax C, Porsius AJ, et al. The completeness of medication histories in hospital medical records of patients admitted to general internal medicine wards. *Br J Clin Pharmacol* 2000;49:597–603.
- Pippins JR, Gandhi TK, Hamann C, et al. Classifying and predicting errors of inpatient medication reconciliation. *J Gen Intern Med* 2008;23:1414–22.
- Lindberg M, Lindberg P, Wikstrom B. Medication discrepancy: a concordance problem between dialysis patients and caregivers. *Scand J Urol Nephrol* 2007;41:546–52.
- Glintborg B, Hillestrom PR, Olsen LH, et al. Are patients reliable when self-reporting medication use? Validation of structured drug interviews and home visits by drug analysis and prescription data in acutely hospitalized patients. *J Clin Pharmacol* 2007;47:1440–9.
- Yang JC, Tomlinson G, Naglie G. Medication lists for elderly patients: clinic-derived versus in-home inspection and interview. *J Gen Intern Med* 2001;16:112–15.
- Gurwitz JH, Field TS, Harrold LR, et al. Incidence and preventability of adverse drug events among older persons in the ambulatory setting. *JAMA* 2003;289:1107–16.
- Piazza G, Nguyen TN, Cios D, et al. Anticoagulation-associated adverse drug events. *Am J Med* 2011;124:1136–42.
- Alam M, Goldberg LH. Serious adverse vascular events associated with perioperative interruption of antiplatelet and anticoagulant therapy. *Dermatol Surg* 2002;28:992–8.
- Ho PM, Peterson ED, Wang L, et al. Incidence of death and acute myocardial infarction associated with stopping clopidogrel after acute coronary syndrome. *JAMA* 2008;299:532–9.

Research and applications

- 31 Choo PW, Rand CS, Inui TS, *et al.* Validation of patient reports, automated pharmacy records, and pill counts with electronic monitoring of adherence to antihypertensive therapy. *Med Care* 1999;37:846–57.
- 32 Lau HS, De BA, Beuning KS, *et al.* Validation of pharmacy records in drug exposure assessment. *J Clin Epidemiol* 1997;50:619–25.
- 33 Bramley TJ, Gerbino PP, Nightengale BS, *et al.* Relationship of blood pressure control to adherence with antihypertensive monotherapy in 13 managed care organizations. *J Manag Care Pharm* 2006;12:239–45.
- 34 Ho PM, Rumsfeld JS, Masoudi FA, *et al.* Effect of medication nonadherence on hospitalization and mortality among patients with diabetes mellitus. *Arch Intern Med* 2006;166:1836–41.
- 35 Stern L, Berman J, Lumry W, *et al.* Medication compliance and disease exacerbation in patients with asthma: a retrospective study of managed care data. *Ann Allergy Asthma Immunol* 2006;97:402–8.
- 36 Sikka R, Xia F, Aubert RE. Estimating medication persistency using administrative claims data. *Am J Manag Care* 2005;11:449–57.
- 37 Steiner JF, Prochazka AV. The assessment of refill compliance using pharmacy records: methods, validity, and applications. *J Clin Epidemiol* 1997;50:105–16.
- 38 Maenpaa T, Suominen T, Asikainen P, *et al.* The outcomes of regional healthcare information systems in health care: a review of the research literature. *Int J Med Inform* 2009;78:757–71.
- 39 Protti D. Comparison of information technology in general practice in 10 countries. *Healthc Q* 2007;10:107–16.
- 40 Elliott RA, Woodward MC. Medication-related problems in patients referred to aged care and memory clinics at a tertiary care hospital. *Australas J Ageing* 2011;30:124–9.
- 41 Lizer MH, Brackbill ML. Medication history reconciliation by pharmacists in an inpatient behavioral health unit. *Am J Health Syst Pharm* 2007;64:1087–91.
- 42 Overhage JM, Dexter PR, Perkins SM, *et al.* A randomized, controlled trial of clinical information shared from another institution. *Ann Emerg Med* 2002;39:14–23.
- 43 Vest JR, Zhao H, Jasperson J, *et al.* Factors motivating and affecting health information exchange usage. *J Am Med Inform Assoc* 2011;18:143–9.
- 44 Adler-Milstein J, Bates DW, Jha AK. A survey of health information exchange organizations in the United States: implications for meaningful use. *Ann Intern Med* 2011;154:666–71.
- 45 Frisse ME, Johnson KB, Nian H, *et al.* The financial impact of health information exchange on emergency department care. *J Am Med Inform Assoc* 2012;19:328–33.
- 46 Lesselroth BJ, Felder RS, Adams SM, *et al.* Design and implementation of a medication reconciliation kiosk: the Automated Patient History Intake Device (APHID). *J Am Med Inform Assoc* 2009;16:300–4.



Estimating the information gap between emergency department records of community medication compared to on-line access to the community-based pharmacy records

Robyn Tamblyn, Lise Poissant, Allen Huang, et al.

J Am Med Inform Assoc 2014 21: 391-398 originally published online August 16, 2013

doi: 10.1136/amiajnl-2013-001704

Updated information and services can be found at:

<http://jamia.bmj.com/content/21/3/391.full.html>

These include:

References

This article cites 45 articles, 10 of which can be accessed free at:

<http://jamia.bmj.com/content/21/3/391.full.html#ref-list-1>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (146 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>